

Recombinant Anti-CD14 Rabbit mAb Product Manual

Cat No: GMA1062

Source: Rabbit

Reactivity: H, M, R

Applications: WB, IH

*Application Key:

H - Human, M - Mouse, R - Rat, B - Bovine, C - Chicken, D - Dog, G - Goat, Mk - Monkey, P - Pig, Rb - Rabbit, S - Sheep, Z - Zebrafish

*Species Reactivity Key:

E- ELISA, WB - Western blot, IH - Immunohistochemistry, IF - Immunofluorescence, FC - Flow cytometry, IC - Immunocytochemistry, IP - Immunoprecipitation, ChIP - Chromatin Immunoprecipitation, EMSA - Electrophoretic Mobility Shift Assay, BL - Blocking, SE - Sandwich ELISA, CBE - Cell-based ELISA, RNAi - RNA interference

Datasheet

Description: Recombinant rabbit monoclonal antibody to CD14

Immunogen: Recombinant fusion protein of human CD14. The exact sequence is proprietary.

Purification: The antibody was purified by immunogen affinity chromatography.

Clonality: Monoclonal

Form: Liquid in PBS, pH 7.3, 50% glycerol, and 0.05% Proclin300.

Dilution: WB (1/500 - 1/1000), IH (1/100 - 1/500)

Gene Symbol: CD14

Alternative Names: Monocyte differentiation antigen CD14; Myeloid cell-specific leucine-rich glycoprotein; CD14

Entrez Gene (Human): 929;

Entrez Gene (Mouse): 12475;

Entrez Gene (Rat): 60350;


SwissProt (Human): P08571;

SwissProt (Mouse): P10810;


SwissProt (Rat): Q63691;

Storage/Stability: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Validation



Western blot analysis of CD14 expression in HepG2 (A), THP1 (B), MCF7 (C), mouse liver (D), rat kidney (E), rat liver (F) whole cell lysates. (Predicted band size: 40 kD; Observed band size: 50-55, 40 kD)



Immunohistochemical analysis of CD14 staining in human colon formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

官方网址: <http://www.genesion.com.cn>

订货热线: 4006169114、020-84224925

Email: whiga22@126.com

