

Recombinant Anti-Desmin Rabbit mAb Product Manual

Cat No: GMA1098

Source: Rabbit

Reactivity: H, M, R

Applications: WB, IH

*Application Key:

H - Human, M - Mouse, R - Rat, B - Bovine, C - Chicken, D - Dog, G - Goat, Mk - Monkey, P - Pig, Rb - Rabbit, S - Sheep, Z - Zebrafish

*Species Reactivity Key:

E- ELISA, WB - Western blot, IH - Immunohistochemistry, IF - Immunofluorescence, FC - Flow cytometry, IC - Immunocytochemistry, IP - Immunoprecipitation, ChIP - Chromatin Immunoprecipitation, EMSA - Electrophoretic Mobility Shift Assay, BL - Blocking, SE - Sandwich ELISA, CBE - Cell-based ELISA, RNAi - RNA interference

Datasheet

Description: Recombinant rabbit monoclonal antibody to Desmin

Immunogen: Recombinant fusion protein of human Desmin. The exact sequence is proprietary.

Purification: The antibody was purified by immunogen affinity chromatography.

Clonality: Monoclonal

Form: Liquid in PBS, pH 7.3, 50% glycerol, and 0.05% Proclin300.

Dilution: WB (1/500 - 1/1000), IH (1/100 - 1/500)

Gene Symbol: DES

Alternative Names: Desmin

Entrez Gene (Human): 1674;

Entrez Gene (Mouse): 13346;

Entrez Gene (Rat): 64362;


SwissProt (Human): P17661;

SwissProt (Mouse): P31001;


SwissProt (Rat): P48675;

Storage/Stability: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Validation



Western blot analysis of Desmin expression in HepG2 (A), MCF7 (B), mouse muscle (C), rat muscle (D) whole cell lysates. (Predicted band size: 53 kD; Observed band size: 45-55 kD)



Immunohistochemical analysis of Desmin staining in human colon formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of Desmin staining in human cardiac muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

官方网站: <http://www.genesion.com.cn>

订货热线: 4006169114, 020-84224925

Email: whiga22@126.com

