

Recombinant Anti-STING Rabbit mAb Product Manual

Cat No: GMA1177

Source: Rabbit

Reactivity: H, M, R

Applications: WB, IH

*Application Key:

H - Human, M - Mouse, R - Rat, B - Bovine, C - Chicken, D - Dog, G - Goat, Mk - Monkey, P - Pig, Rb - Rabbit, S - Sheep, Z - Zebrafish

*Species Reactivity Key:

E - ELISA, WB - Western blot, IH - Immunohistochemistry, IF - Immunofluorescence, FC - Flow cytometry, IC - Immunocytochemistry, IP - Immunoprecipitation, ChIP - Chromatin Immunoprecipitation, EMSA - Electrophoretic Mobility Shift Assay, BL - Blocking, SE - Sandwich ELISA, CBE - Cell-based ELISA, RNAi - RNA interference

Datasheet

Description: Recombinant rabbit monoclonal antibody to STING

Immunogen: Recombinant fusion protein of human STING. The exact sequence is proprietary.

Purification: The antibody was purified by immunogen affinity chromatography.

Clonality: Monoclonal

Form: Liquid in PBS, pH 7.3, 50% glycerol, and 0.05% Proclin300.

Dilution: WB (1/500 - 1/1000), IH (1/100 - 1/500)

Gene Symbol: STING1

Alternative Names: ERIS; MITA; STING; TMEM173; Stimulator of interferon genes protein; hSTING; Endoplasmic reticulum interferon stimulator; ERIS; Mediator of IRF3 activation; hMITA; Transmembrane protein 173

Entrez Gene (Human): 340061;


Entrez Gene (Mouse): 72512;

SwissProt (Human): Q86WV6;


SwissProt (Mouse): Q3TBT3;

Storage/Stability: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Validation



Western blot analysis of STING expression in MG63 (A), HUT78 (B), mouse kidney (C), mouse lung (D), rat kidney (E), rat lung (F) whole cell lysates. (Predicted band size: 42 kD; Observed band size: 35 kD)



Immunohistochemical analysis of STING staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

官方网址: <http://www.genscion.com.cn>

订货热线: 4006169114, 020-84224925

Email: whiga22@126.com

