

Anti-PCNA Antibody [KO/KD Validated] Product Manual

Cat No: GPA1859

Source: Rabbit

Reactivity: H, M, R, B, C, D, Mk, P, Rb

Applications: WB, IH

*Application Key:

H - Human, M - Mouse, R - Rat, B - Bovine, C - Chicken, D - Dog, G - Goat, Mk - Monkey, P - Pig, Rb - Rabbit, S - Sheep, Z - Zebrafish

*Species Reactivity Key:

E- ELISA, WB - Western blot, IH - Immunohistochemistry, IF - Immunofluorescence, FC - Flow cytometry, IC - Immunocytochemistry, IP - Immunoprecipitation, ChIP - Chromatin Immunoprecipitation, EMSA - Electrophoretic Mobility Shift Assay, BL - Blocking, SE - Sandwich ELISA, CBE - Cell-based ELISA, RNAi - RNA interference

Datasheet

Description: Rabbit polyclonal antibody to PCNA

Immunogen: KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PCNA. The exact sequence is proprietary.

Purification: The antibody was purified by immunogen affinity chromatography.

Clonality: Polyclonal

Form: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Dilution: WB (1/500 - 1/1000), IH (1/100 - 1/200)

Gene Symbol: PCNA

Alternative Names: Proliferating cell nuclear antigen; PCNA; Cyclin

Entrez Gene (Human): 5111;

Entrez Gene (Mouse): 18538;

Entrez Gene (Rat): 25737;


SwissProt (Human): P12004;

SwissProt (Mouse): P17918;


SwissProt (Rat): P04961;

Storage/Stability: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Validation



Western blot analysis of PCNA expression in H9C2 (A), 3T3L1 (B), HEK293T (C), A549 (D), HCT116 (E) whole cell lysates. (Predicted band size: 28 kD; Observed band size: 36 kD))



Immunohistochemical analysis of PCNA staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blot analysis of PCNA expression in wild type (WT) and knockdown (KD) HeLa cell lysates.

