

Recombinant Anti-Histone H3 (Lactyl-K18) Rabbit mAb

Product Manual

货号: GMA1203

来源: Rabbit

反应物种: H, M, R

实验应用: WB, IH, IF/IC

***反应物种注解:**

H - Human, M - Mouse, R - Rat, B - Bovine, C - Chicken, D - Dog, G - Goat, Mk - Monkey, P - Pig, Rb - Rabbit, S - Sheep, Z - Zebrafish

***实验应用注解:**

E - ELISA, WB - Western blot, IH - Immunohistochemistry, IF - Immunofluorescence, FC - Flow cytometry, IC - Immunocytochemistry, IP - Immunoprecipitation, ChIP - Chromatin Immunoprecipitation, EMSA - Electrophoretic Mobility Shift Assay, BL - Blocking, SE - Sandwich ELISA, CBE - Cell-based ELISA, RNAi - RNA interference

Datasheet

产品描述: Recombinant rabbit monoclonal antibody to Histone H3 (Lactyl-K18)

免疫原: KLH-conjugated synthetic lactylated peptide corresponding to residues surrounding K18 of human Histone H3 protein.

The exact sequence is proprietary.

纯化方式: This antibody is purified through a protein G column.

克隆类型: Monoclonal

产品形式: Liquid in PBS, pH 7.3, 50% glycerol, and 0.05% Proclin300.

稀释比: WB (1/1000 - 1/2000), IH (1/50 - 1/200), IF/IC (1/50 - 1/200)

基因名称: HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J;

HIST2H3A; HIST2H3C; HIST2H3D; H3F3A; H3F3B

相关名称: HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/h; Histone H3/i; Histone H3/j; Histone H3/k; Histone H3/l; HIST2H3A; HIST2H3C; H3F2; H3FM;

HIST2H3D; Histone H3.2; Histone H3/m; Histone H3/o; H3F3A; H3.3A; H3F3; PP781; H3F3B; H3.3B; Histone H3.3

基因编号(人): 8350; 8351; 8352; 8353; 8354; 8355; 8356; 8357; 8358; 8968; 126961; 333932; 653604; 3020; 3021;

基因编号(小鼠): 319152; 15077; 15078;

基因编号(大鼠): 291159; 100361558; 蛋白编号


(人): P68431; Q71DI3; P84243; 蛋白编号(小鼠):

P68433; P84228; P84244;


蛋白编号(大鼠): Q6LED0; P84245;

储存效期: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Validation




Western blot analysis of Histone H3 (Lactyl-K18) expression in Jurkat (A), mouse liver (B), mouse muscle (C), rat liver (D), rat muscle (E) whole cell lysates. (Predicted band size: 15 kD; Observed band size: 17 kD)



Immunohistochemical analysis of Histone H3 (Lactyl-K18) staining in human stomach cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Histone H3 (Lactyl-K18) staining in 22RV1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



Direct ELISA antibody dose-response curve using Recombinant Anti-Histone H3 (Lactyl-K18) Rabbit mAb. Antigen (lactylated peptide and non-lactylated-peptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

官方网址: <http://www.genscion.com.cn>

订货热线: 4006169114、020-84224925

Email: whiga22@126.com

